

# Additional pages for your INQOVI AML Health Journal



These pages are designed to help you continue using the INQOVI Health Journal you received in your Patient Welcome Kit. You've already been tracking your treatment and side effects—this refill is here to support you as you move forward.

Inside, you'll find **6 sets of pages**, with each set containing a **side effect tracker** and a **treatment tracker** to keep recording how you're feeling and your daily doses. There is also **one notes page** at the end for anything else you'd like to remember or share with your healthcare team.

If you need more pages along the way, you can reprint this refill anytime to support you throughout your INQOVI treatment journey.

**See the brochure in your INQOVI Patient Welcome Kit for helpful information on taking and tracking your INQOVI, possible side effects, and support and resources.**

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

**INQOVI**<sup>®</sup>  
(decitabine and cedazuridine)  
35mg / 100mg tablets

## INQOVI AML Side Effect Tracker

If you experience any side effects while taking INQOVI, you can mark the severity and include related notes below to discuss them with your doctor. Not all side effects are shown here. Some may only show up in blood tests your doctor orders. For more information on possible side effects, see the brochure in your INQOVI Patient Welcome Kit.

Visit [INQOVI.com](http://INQOVI.com) for more information, and talk to your healthcare provider if you have any questions or concerns.

Month \_\_\_\_\_

SIDE EFFECT	None	Mild	Moderate	Severe	Date(s) experienced/Notes
Arm or leg swelling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Bleeding	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Constipation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Decreased appetite	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Diarrhea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Dizziness or feeling lightheaded	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Joint pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Mouth or throat pain or sores	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Muscle pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Nausea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Rash	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Shortness of breath	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Stomach-area (abdomen) pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Tiredness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Other	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

## INQOVI AML Treatment Tracker



Keep following this schedule for as long as your doctor tells you to. If your treatment does not follow this schedule, ask your doctor to note where it is different.

To find out more about taking venetoclax, please see venetoclax Prescribing Information.

For all cycles, remember to:

- ✓ Take INQOVI at approximately the same time each day
- ✓ Take INQOVI on an empty stomach at least 2 hours before or 2 hours after eating
- ✓ Swallow the tablet whole, without cutting, crushing, or chewing it
- ✓ Take INQOVI and venetoclax at least 2 hours apart. **It's important that you do not take INQOVI and venetoclax at the same time**

Month \_\_\_\_\_

ADDITIONAL CYCLES (This is for any cycle after your first INQOVI cycle)						
Day 1 	Day 2 	Day 3 	Day 4 	Day 5 	Day 6 	Day 7 
Day 8 	Day 9 	Day 10 	Day 11 	Day 12 	Day 13 	Day 14 
Day 15 	Day 16 	Day 17 	Day 18 	Day 19 	Day 20 	Day 21 
Day 22 	Day 23 	Day 24 	Day 25 	Day 26 	Day 27 	Day 28 

INQOVI = 

4 pills of venetoclax = 

Venetoclax is for patients with AML only.

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

**INQOVI**<sup>®</sup>  
(decitabine and cedazuridine)  
35mg / 100mg tablets

## INQOVI AML Side Effect Tracker

If you experience any side effects while taking INQOVI, you can mark the severity and include related notes below to discuss them with your doctor. Not all side effects are shown here. Some may only show up in blood tests your doctor orders. For more information on possible side effects, see the brochure in your INQOVI Patient Welcome Kit.

Visit [INQOVI.com](http://INQOVI.com) for more information, and talk to your healthcare provider if you have any questions or concerns.

Month \_\_\_\_\_

SIDE EFFECT	None	Mild	Moderate	Severe	Date(s) experienced/Notes
Arm or leg swelling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Bleeding	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Constipation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Decreased appetite	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Diarrhea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Dizziness or feeling lightheaded	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Joint pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Mouth or throat pain or sores	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Muscle pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Nausea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Rash	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Shortness of breath	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Stomach-area (abdomen) pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Tiredness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Other	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

## INQOVI AML Treatment Tracker




Keep following this schedule for as long as your doctor tells you to. If your treatment does not follow this schedule, ask your doctor to note where it is different.

To find out more about taking venetoclax, please see venetoclax Prescribing Information.

For all cycles, remember to:

- ✓ Take INQOVI at approximately the same time each day
- ✓ Take INQOVI on an empty stomach at least 2 hours before or 2 hours after eating
- ✓ Swallow the tablet whole, without cutting, crushing, or chewing it
- ✓ Take INQOVI and venetoclax at least 2 hours apart. **It's important that you do not take INQOVI and venetoclax at the same time**

Month \_\_\_\_\_

ADDITIONAL CYCLES (This is for any cycle after your first INQOVI cycle)						
Day 1 	Day 2 	Day 3 	Day 4 	Day 5 	Day 6 	Day 7 
Day 8 	Day 9 	Day 10 	Day 11 	Day 12 	Day 13 	Day 14 
Day 15 	Day 16 	Day 17 	Day 18 	Day 19 	Day 20 	Day 21 
Day 22 	Day 23 	Day 24 	Day 25 	Day 26 	Day 27 	Day 28 

INQOVI = 

4 pills of venetoclax = 

Venetoclax is for patients with AML only.

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

**INQOVI**<sup>®</sup>  
(decitabine and cedazuridine)  
35mg / 100mg tablets

## INQOVI AML Side Effect Tracker

If you experience any side effects while taking INQOVI, you can mark the severity and include related notes below to discuss them with your doctor. Not all side effects are shown here. Some may only show up in blood tests your doctor orders. For more information on possible side effects, see the brochure in your INQOVI Patient Welcome Kit.

Visit [INQOVI.com](http://INQOVI.com) for more information, and talk to your healthcare provider if you have any questions or concerns.

Month \_\_\_\_\_

SIDE EFFECT	None	Mild	Moderate	Severe	Date(s) experienced/Notes
Arm or leg swelling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Bleeding	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Constipation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Decreased appetite	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Diarrhea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Dizziness or feeling lightheaded	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Joint pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Mouth or throat pain or sores	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Muscle pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Nausea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Rash	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Shortness of breath	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Stomach-area (abdomen) pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Tiredness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Other	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

## INQOVI AML Treatment Tracker

Keep following this schedule for as long as your doctor tells you to. If your treatment does not follow this schedule, ask your doctor to note where it is different.

To find out more about taking venetoclax, please see venetoclax Prescribing Information.

For all cycles, remember to:

- ✓ Take INQOVI at approximately the same time each day
- ✓ Take INQOVI on an empty stomach at least 2 hours before or 2 hours after eating
- ✓ Swallow the tablet whole, without cutting, crushing, or chewing it
- ✓ Take INQOVI and venetoclax at least 2 hours apart. **It's important that you do not take INQOVI and venetoclax at the same time**

Month \_\_\_\_\_

ADDITIONAL CYCLES (This is for any cycle after your first INQOVI cycle)						
Day 1 	Day 2 	Day 3 	Day 4 	Day 5 	Day 6 	Day 7 
Day 8 	Day 9 	Day 10 	Day 11 	Day 12 	Day 13 	Day 14 
Day 15 	Day 16 	Day 17 	Day 18 	Day 19 	Day 20 	Day 21 
Day 22 	Day 23 	Day 24 	Day 25 	Day 26 	Day 27 	Day 28 

INQOVI = 

4 pills of venetoclax = 

Venetoclax is for patients with AML only.

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

**INQOVI**<sup>®</sup>  
(decitabine and cedazuridine)  
35mg / 100mg tablets

## INQOVI AML Side Effect Tracker

If you experience any side effects while taking INQOVI, you can mark the severity and include related notes below to discuss them with your doctor. Not all side effects are shown here. Some may only show up in blood tests your doctor orders. For more information on possible side effects, see the brochure in your INQOVI Patient Welcome Kit.

Visit [INQOVI.com](http://INQOVI.com) for more information, and talk to your healthcare provider if you have any questions or concerns.

Month \_\_\_\_\_

SIDE EFFECT	None	Mild	Moderate	Severe	Date(s) experienced/Notes
Arm or leg swelling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Bleeding	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Constipation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Decreased appetite	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Diarrhea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Dizziness or feeling lightheaded	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Joint pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Mouth or throat pain or sores	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Muscle pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Nausea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Rash	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Shortness of breath	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Stomach-area (abdomen) pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Tiredness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Other	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

## INQOVI AML Treatment Tracker

Keep following this schedule for as long as your doctor tells you to. If your treatment does not follow this schedule, ask your doctor to note where it is different.

To find out more about taking venetoclax, please see venetoclax Prescribing Information.

For all cycles, remember to:

- ✓ Take INQOVI at approximately the same time each day
- ✓ Take INQOVI on an empty stomach at least 2 hours before or 2 hours after eating
- ✓ Swallow the tablet whole, without cutting, crushing, or chewing it
- ✓ Take INQOVI and venetoclax at least 2 hours apart. **It's important that you do not take INQOVI and venetoclax at the same time**

Month \_\_\_\_\_

ADDITIONAL CYCLES (This is for any cycle after your first INQOVI cycle)						
Day 1 	Day 2 	Day 3 	Day 4 	Day 5 	Day 6 	Day 7 
Day 8 	Day 9 	Day 10 	Day 11 	Day 12 	Day 13 	Day 14 
Day 15 	Day 16 	Day 17 	Day 18 	Day 19 	Day 20 	Day 21 
Day 22 	Day 23 	Day 24 	Day 25 	Day 26 	Day 27 	Day 28 

INQOVI = 

4 pills of venetoclax = 

Venetoclax is for patients with AML only.

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

**INQOVI**<sup>®</sup>  
(decitabine and cedazuridine)  
35mg / 100mg tablets

## INQOVI AML Side Effect Tracker

If you experience any side effects while taking INQOVI, you can mark the severity and include related notes below to discuss them with your doctor. Not all side effects are shown here. Some may only show up in blood tests your doctor orders. For more information on possible side effects, see the brochure in your INQOVI Patient Welcome Kit.

Visit [INQOVI.com](http://INQOVI.com) for more information, and talk to your healthcare provider if you have any questions or concerns.

Month \_\_\_\_\_

SIDE EFFECT	None	Mild	Moderate	Severe	Date(s) experienced/Notes
Arm or leg swelling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Bleeding	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Constipation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Decreased appetite	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Diarrhea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Dizziness or feeling lightheaded	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Joint pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Mouth or throat pain or sores	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Muscle pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Nausea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Rash	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Shortness of breath	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Stomach-area (abdomen) pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Tiredness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Other	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

## INQOVI AML Treatment Tracker




Keep following this schedule for as long as your doctor tells you to. If your treatment does not follow this schedule, ask your doctor to note where it is different.

To find out more about taking venetoclax, please see venetoclax Prescribing Information.

For all cycles, remember to:

- Take INQOVI at approximately the same time each day
- Take INQOVI on an empty stomach at least 2 hours before or 2 hours after eating
- Swallow the tablet whole, without cutting, crushing, or chewing it
- Take INQOVI and venetoclax at least 2 hours apart. **It's important that you do not take INQOVI and venetoclax at the same time**

Month \_\_\_\_\_

ADDITIONAL CYCLES (This is for any cycle after your first INQOVI cycle)						
Day 1  	Day 2  	Day 3  	Day 4  	Day 5  	Day 6 	Day 7 
Day 8 	Day 9 	Day 10 	Day 11 	Day 12 	Day 13 	Day 14 
Day 15 	Day 16 	Day 17 	Day 18 	Day 19 	Day 20 	Day 21 
Day 22 	Day 23 	Day 24 	Day 25 	Day 26 	Day 27 	Day 28 

INQOVI = 

4 pills of venetoclax = 

Venetoclax is for patients with AML only.

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

**INQOVI**<sup>®</sup>  
(decitabine and cedazuridine)  
35mg / 100mg tablets

## INQOVI AML Side Effect Tracker

If you experience any side effects while taking INQOVI, you can mark the severity and include related notes below to discuss them with your doctor. Not all side effects are shown here. Some may only show up in blood tests your doctor orders. For more information on possible side effects, see the brochure in your INQOVI Patient Welcome Kit.

Visit [INQOVI.com](http://INQOVI.com) for more information, and talk to your healthcare provider if you have any questions or concerns.

Month \_\_\_\_\_

SIDE EFFECT	None	Mild	Moderate	Severe	Date(s) experienced/Notes
Arm or leg swelling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Bleeding	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Constipation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Decreased appetite	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Diarrhea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Dizziness or feeling lightheaded	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Joint pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Mouth or throat pain or sores	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Muscle pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Nausea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Rash	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Shortness of breath	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Stomach-area (abdomen) pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Tiredness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	
Other	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

## INQOVI AML Treatment Tracker

Keep following this schedule for as long as your doctor tells you to. If your treatment does not follow this schedule, ask your doctor to note where it is different.

To find out more about taking venetoclax, please see venetoclax Prescribing Information.

For all cycles, remember to:

- ✓ Take INQOVI at approximately the same time each day
- ✓ Take INQOVI on an empty stomach at least 2 hours before or 2 hours after eating
- ✓ Swallow the tablet whole, without cutting, crushing, or chewing it
- ✓ Take INQOVI and venetoclax at least 2 hours apart. **It's important that you do not take INQOVI and venetoclax at the same time**

Month \_\_\_\_\_

ADDITIONAL CYCLES (This is for any cycle after your first INQOVI cycle)						
Day 1 	Day 2 	Day 3 	Day 4 	Day 5 	Day 6 	Day 7 
Day 8 	Day 9 	Day 10 	Day 11 	Day 12 	Day 13 	Day 14 
Day 15 	Day 16 	Day 17 	Day 18 	Day 19 	Day 20 	Day 21 
Day 22 	Day 23 	Day 24 	Day 25 	Day 26 	Day 27 	Day 28 

INQOVI = 

4 pills of venetoclax = 

Venetoclax is for patients with AML only.

This is not meant to replace advice given to you by a doctor. Always contact your healthcare team if you have questions or concerns with taking your medication as instructed.

**INQOVI**<sup>®</sup>  
(decitabine and cedazuridine)  
35mg / 100mg tablets



## What is INQOVI?

INQOVI is a prescription medicine used:

- to treat adults with myelodysplastic syndromes (MDS), including chronic myelomonocytic leukemia (CMML). Your healthcare provider will determine if INQOVI can treat your type of MDS
- in combination with venetoclax to treat adults with newly diagnosed acute myeloid leukemia (AML) who:
  - are 75 years of age or older, or
  - have other medical conditions that prevent the use of standard chemotherapy

It is not known if INQOVI is safe or effective in children.

## Important Safety Information

**INQOVI may cause serious side effects, including:**

**Low blood cell counts.** Low blood counts (white blood cells, platelets, and red blood cells) are common with INQOVI but can also be serious and lead to infections that may be life-threatening. If your blood cell counts are too low, your healthcare provider may need to delay treatment with INQOVI, lower your dose of INQOVI, or in some cases, give you a medicine to help treat low blood cell counts. Your healthcare provider may need to give you antibiotic medicines to prevent or treat infections or fever while your blood cell counts are low. Your healthcare provider will check your blood cell counts before you start treatment and regularly during treatment with INQOVI.

**Tell your healthcare provider right away if you get any of the following signs and symptoms during treatment with INQOVI:**

- fever
- chills
- body aches
- tiredness
- bruising more easily than usual

**Before taking INQOVI, tell your healthcare provider about all of your medical conditions, including if you:**

- have kidney problems
- are pregnant or plan to become pregnant. INQOVI can harm your unborn baby

**Females who are able to become pregnant:**

- Your healthcare provider will check to see if you are pregnant before you start treatment with INQOVI
- Use effective birth control during treatment with INQOVI and for 6 months after the last dose of INQOVI

**Males with female partners who are able to become pregnant** should use effective birth control during treatment with INQOVI and for 3 months after the last dose. Talk to your healthcare provider if you have questions about birth control options that are right for you.

- are breastfeeding or plan to breastfeed. It is not known if INQOVI passes into breast milk. **Do not** breastfeed during treatment with INQOVI and for 2 weeks after your last dose of INQOVI

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. INQOVI may affect the way certain other medicines work and may cause side effects. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

**If taking INQOVI in combination with venetoclax for AML, please refer to the venetoclax Medication Guide for additional information.**

**The most common side effects of INQOVI in adults with MDS or CMML include:** low white blood cell count (leukopenia, neutropenia), low platelets in your blood (thrombocytopenia), low red blood cell count (anemia), tiredness, constipation, bleeding, muscle pain, pain or sores in your mouth or throat, joint pain, nausea, shortness of breath, diarrhea, rash, dizziness, fever with low white blood cell count (febrile neutropenia), swelling of arms or legs, headache, cough, decreased appetite, upper respiratory tract infection, pneumonia, and changes in liver function tests.

**The most common side effects of INQOVI in adults with AML in combination with venetoclax include:** low white blood cell counts (leukopenia, lymphocytopenia, neutropenia), low platelets in your blood (thrombocytopenia), fever with low white blood cell count (febrile neutropenia), bleeding, low red blood cell count (anemia), infection, diarrhea, tiredness, pain or sores in your mouth or throat, constipation, joint pain, decreased appetite, swelling of arms or legs, nausea, shortness of breath, sepsis, pneumonia, rash, changes in liver function tests, muscle pain, abnormal heart rhythm, stomach-area (abdomen) pain, dizziness or feeling lightheaded, and decreased kidney function.

INQOVI may affect fertility in men. Talk to your healthcare provider if this is a concern for you. These are not all of the possible side effects of INQOVI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**If you are taking INQOVI in combination with venetoclax for AML, please refer to the venetoclax Medication Guide for additional information.**

**Please see full Prescribing Information at [INQOVI.com/PI](http://INQOVI.com/PI), and discuss with your healthcare provider.**

Marketed by © Taiho Oncology, Inc. INQOVI is a registered trademark of Taiho Pharmaceutical Co., Ltd., used under license by Taiho Oncology, Inc. © TAIHO ONCOLOGY, INC. 101 Carnegie Center, Suite 300, Princeton, NJ 08540. All rights reserved.

 TAIHO ONCOLOGY

© TAIHO ONCOLOGY, INC. 05/2026  
INQ-PM-US-0623

**INQOVI**<sup>®</sup>  
(decitabine and cedazuridine)  
35mg / 100mg tablets